

PERIODONTAL CHALLENGES IN CERAMIC VENEERING: MANAGEMENT OF EMERGENCE PROFILE AND SOFT TISSUES

Diana Hanu¹, Dan Zagnat^{1*}, Forna Norina²

1 DDS, Resident, Faculty of Dental Medicine, University of Medicine and Pharmacy "Grigore T.Popa", Iasi

2 "Gr. T. Popa" U.M.Ph. - Iași, Romania, Faculty of Dentistry, Department of Implantology, Removable Dentures, Dentures Technology

Corresponding authors: Dan Zagnat , e-mail: dzagnat@gmail.com

ABSTRACT

Aim of the study The objective of this review is to evaluate contemporary clinical protocols for the management of emergence profiles and peri-veneer soft tissues. The study aims to highlight the critical necessity of a multidisciplinary approach where periodontal stability serves as the foundation for the long-term success of ceramic veneer therapy. **Material and methods** The review synthesizes current literature regarding the interaction between restorative design and periodontal biology. Key parameters analyzed include the preservation of biological width, the impact of ceramic micro-topography on cell adhesion, the management of residual cement, the diagnostic significance of the periodontal biotype, and the precise control of the emergence profile and gingival zenith. **Conclusions** Clinicians must balance patient aesthetic expectations with the biological limits of the periodontal apparatus. By integrating meticulous soft tissue management, standardized polishing protocols, and rigorous long-term monitoring, practitioners can achieve a stable, biocompatible, and high-level aesthetic integration in ceramic veneer therapy.

Key words: ceramic veneers, biological width, emergence profile, periodontal biotype, gingival zenith.

1. INTRODUCTION

In the contemporary landscape of restorative dentistry, the pursuit of aesthetic perfection has shifted from simple tooth modification to a complex biological integration. Ceramic veneers represent one of the most minimally invasive and aesthetically rewarding treatments; however, their long-term success is fundamentally anchored in periodontal health (Al-Shorman *et al.*, 2024). As clinicians, we must recognize that the transition zone between the restorative material and the periodontal tissues constitutes a critical interface that dictates the longevity and gingival harmony of the final result.

The management of the emergence profile and the surrounding soft tissues represents a significant clinical challenge, particularly in patients with delicate periodontal biotypes (Zihan *et al.*, 2025). Improper contouring or violation of the biological width during preparation can lead

to chronic inflammation, gingival recession, and the loss of interdental papillae, effectively compromising the aesthetic outcome (Karvekar *et al.*, 2025). Therefore, achieving an optimal aesthetic result requires a comprehensive understanding of the interplay between restorative design and periodontal response. This review aims to examine the current clinical protocols for managing soft tissues, emphasizing the necessity of a multidisciplinary approach where periodontal stability serves as the foundation for successful ceramic veneer therapy.

The increasing demand for esthetic rehabilitation in patients with pre-existing periodontal conditions necessitates a paradigm shift in treatment planning. Historically, the focus was primarily on the mechanical retention and optical properties of the ceramic material. However, modern clinical practice highlights the imperative of prioritizing periodontal biology, as any

restorative procedure that ignores the gingival architecture is inherently prone to failure. Consequently, the clinician must balance the patient's esthetic expectations with the biological limits of the periodontal apparatus, ensuring that restorative margins facilitate proper plaque control and support optimal gingival health (Ercoli *et al.*, 2021).

Furthermore, the integration of digital technology and advanced surgical techniques has significantly expanded our ability to manipulate soft tissue contours. By utilizing provisional restorations to define the desired emergence profile, practitioners can guide the maturation of the gingival margin prior to the placement of the final ceramic veneers (Andrei *et al.*, 2024). This proactive approach not only minimizes the risk of post-operative complications but also serves as a diagnostic tool for assessing long-term tissue stability.

2. MATERIAL AND METHODS

2.1. Biological Width

In the context of ceramic veneer restorations, the biological width represents a fundamental concept for maintaining periodontal health and ensuring the long-term success of the treatment (Talpos *et al.*, 2025). It is defined as the vertical dimension of the complex composed of the junctional epithelium and the supracrestal connective tissue attachment.

According to the classic studies by Gargiulo *et al.* (1961), the biological width occupies, on average, 2.04 mm in the corono-apical direction, consisting of:

- Gingival sulcus: approximately 0.69 mm.
- Junctional epithelium: approximately 0.97 mm (epithelial attachment).
- Connective tissue attachment: approximately 1.07 mm (fibers inserted into the root cementum).

This anatomical entity acts as a natural biological barrier, protecting the underlying alveolar bone from microbial invasion and mechanical trauma.

When the margin of a ceramic veneer is placed subgingivally, there is an inherent risk of invading the biological width if the preparation finish line is positioned too far apically. This invasion triggers a chronic

inflammatory response, which may progress to:

- ⇒ Gingival recession: loss of epithelial attachment and apical migration of the free gingival margin, compromising esthetics (Niemczyk *et al.*, 2024).
- ⇒ Persistent gingival inflammation: erythema, edema, and bleeding on probing, caused by the tissue's inability to establish a stable attachment on the ceramic surface in close proximity to the alveolar bone (Ercoli *et al.*, 2021).
- ⇒ Loss of attachment: in severe cases, chronic inflammation may lead to alveolar bone resorption as the body attempts to "re-establish" the biological width through apical migration of the attachments, a phenomenon known as bone sounding (Srimaneepong *et al.*, 2022).

To prevent periodontal pathology associated with ceramic veneers, clinical management must include:

- Evaluation of gingival margin position: establishing the position of the finish line (supragingival, equigingival, subgingival) based on esthetic requirements and the patient's periodontal status. Supragingival placement remains the "gold standard" for periodontal health (Cortés-Sánchez *et al.*, 2026).
- Identification of bone level: using bone sounding under local anesthesia to determine the exact distance from the gingival margin to the alveolar crest prior to preparation (De Angelis *et al.*, 2025).
- Respecting depth limits: in cases where subgingival positioning is necessary (for masking tooth discoloration or clinical crown lengthening), it should not exceed 0.5 mm below the gingival sulcus, maintaining a safety margin of at least 2.5 - 3 mm from the alveolar crest (Ceravolo *et al.*, 2026).
- Control of the emergence profile: the transition between the root surface and the ceramic veneer should be as flat as possible (minimal convexity) to prevent plaque accumulation and facilitate the patient's oral hygiene maintenance (Nenov *et al.*, 2025).

2.2. Micro-topography of the ceramic

surface and cell adhesion

In the quest for long-term periodontal stability in prosthetic dentistry, the interaction between the soft tissue complex and the restorative material is of critical importance. While the biological width defines the spatial requirements, the micro-topography of the ceramic surface dictates the quality of the biological seal. The ability of gingival fibroblasts and epithelial cells to adhere, proliferate, and establish a hemidesmosomal attachment is profoundly influenced by the surface characteristics of the restorative material (Paul *et al.*, 2021).

The surface texture of ceramic restorations serves as the interface for the transmucosal attachment (Metwally *et al.*, 2025). Research indicates that surface topography modulates cell behavior through a phenomenon known as contact guidance:

- ⇒ Smooth surfaces ($R_a < 0.2 \mu\text{m}$): surfaces polished to a high luster are generally conducive to soft tissue health. They minimize the surface area available for bacterial adhesion and biofilm maturation, thereby reducing the inflammatory challenge to the peri-implant or periodontal tissues.
- ⇒ Rough surfaces ($R_a > 0.2 \mu\text{m}$): increased surface roughness, often resulting from inadequate polishing or surface texturing, promotes greater bacterial colonization. Furthermore, while rougher surfaces may potentially increase the surface area for protein adsorption, in the context of marginal tissues, they often lead to an accumulation of inflammatory mediators that inhibit stable hemidesmosomal attachment.

The chemical and structural composition of the ceramic material interacts synergistically with its micro-topography to determine cellular adhesion efficiency:

- Zirconia (Y-TZP): due to its high density and low affinity for bacterial colonization, zirconia is widely considered the most biocompatible ceramic for transmucosal components (Yinuo *et al.*, 2024). Studies have

demonstrated that gingival fibroblasts exhibit higher proliferation rates and increased expression of adhesion proteins when cultured on polished zirconia compared to titanium or glass-ceramics.

- Lithium Disilicate and Feldspathic Ceramics: these materials, while highly esthetic, require meticulous surface treatment (Payaminia *et al.*, 2021). The glazing process or mechanical polishing is mandatory to achieve the necessary surface energy that prevents the formation of a pathogenic biofilm at the restoration-gingiva interface.

To optimize cell adhesion and ensure the success of the periodontal-restorative interface, clinicians must adhere to stringent protocols:

1. Polishing protocols: mechanical polishing with diamond-impregnated rubbers followed by a final glaze or high-luster polish is essential to reduce surface roughness below the critical threshold of $0.2 \mu\text{m}$.
2. Biofilm management: since surface micro-topography cannot completely eliminate bacterial adhesion, patient-specific oral hygiene instructions remain the primary factor in preventing the inflammatory degradation of the attachment.
3. Prevention of surface damage: clinicians must avoid the use of rough burs or abrasive instruments on the cervical margin of the veneer during cementation or occlusal adjustment, as these create microscopic "nooks" that facilitate microbial colonization and potential gingival recession.

The ultimate objective of managing ceramic micro-topography is to facilitate the formation of a biological seal. When the surface is appropriately finished, the gingival epithelium can form a continuous hemidesmosomal attachment to the ceramic, similar to the attachment to natural tooth structure. This seal acts as a dynamic barrier, physically excluding plaque-associated endotoxins from the underlying connective

tissue, thereby preventing the collapse of the biological width.

2.3. Residual cement control

The clinical success of ceramic veneers is linked to the meticulous removal of excess luting cement at the restoration-gingival margin interface. Residual cement, often overlooked due to its proximity to the soft tissue, acts as a potent iatrogenic factor capable of inducing localized inflammation, leading to a rapid breakdown of the periodontal attachment (Montevecchi *et al.*, 2025).

The presence of subgingival cement remnants serves as a reservoir for bacterial colonization, significantly altering the peri-veneer environment:

- Bacterial reservoirs: the porous surface of residual resin cements provides an ideal substrate for the adhesion of periodontal pathogens. Unlike natural tooth structure, these materials are susceptible to degradation, which can further increase their surface roughness over time, promoting biofilm retention (Skevi *et al.*, 2021).
- Chemical irritation: unpolymerized monomers within the excess cement can leach into the gingival sulcus, triggering a hypersensitivity response or chronic inflammation of the soft tissues, characterized by edema and gingival enlargement (Alharbi *et al.*, 2024).
- Mechanical trauma: the hardened cement acts as a mechanical barrier that prevents the gingival tissue from adapting closely to the restoration surface, thereby impeding the formation of a biological seal.

Detecting residual cement is frequently complicated by the optical properties of modern restorative materials. Many resin cements exhibit limited radiopacity, making their identification on standard periapical radiographs difficult. This creates a "hidden" risk factor, where cement remains undetected until clinical signs of periodontal disease, such as suppuration, persistent bleeding on probing, or localized recession, become apparent. Shade-matching cements, intended to improve the esthetic integration of the veneer, often blend in with the tooth structure

or the restoration itself, making visual detection during the cleaning phase exceptionally challenging.

To mitigate the risks associated with residual cement, clinicians must implement standardized, highly controlled protocols. Utilizing a retraction cord or an alternative gingival displacement technique is essential to allow for full visual access to the margins, ensuring that excess cement can be thoroughly removed without traumatizing the gingival attachment. Applying the cement strategically and using a "tack-cure" technique allows the cement to achieve a gel-like state. At this stage, the excess can be meticulously removed with an explorer or specialized scaler before final polymerization, significantly reducing the risk of leaving remnants.

A thorough assessment of the cervical area using magnified visualization (dental loupes or a surgical microscope) and dental floss is mandatory. The floss should slide smoothly through the interproximal contact without fraying, which would indicate the presence of subgingival cement "tags." After curing, the use of fine-grit finishing strips and polishing cups is required to ensure that the transition between the veneer margin and the tooth structure is smooth and plaque-resistant, effectively finalizing the management of the emergence profile.

Chronic inflammation resulting from residual cement inevitably leads to the migration of the junctional epithelium and the loss of connective tissue attachment, effectively reversing the benefits of a well-designed restoration. Therefore, the control of residual cement must be treated as a critical, non-negotiable step in the delivery of ceramic veneer therapy.

2.4. Periodontal biotype as a prognostic factor

In the interdisciplinary management of ceramic veneers, the periodontal biotype (or periodontal phenotype) is a critical diagnostic parameter that dictates the therapeutic strategy and influences the long-term clinical prognosis (Parihar *et al.*, 2025). Defined by the morphological characteristics of the

gingival tissues, this factor serves as a primary determinant of how the periodontium will react to the surgical and restorative phases of veneer treatment.

Clinical observation and standardized diagnostic procedures generally categorize the periodontal biotype into two primary phenotypes, each presenting distinct risks for ceramic veneer therapy: thin-scalloped phenotype (characterized by a delicate, translucent gingival architecture with minimal keratinized tissue and a underlying thin alveolar bone plate; this phenotype is highly susceptible to mechanical trauma and surgical insult, with a high propensity for recession following even minor iatrogenic irritation or aggressive preparation) and thick-flat phenotype (distinguished by dense, fibrotic gingival tissue and a more robust alveolar bone morphology; this phenotype is more resilient to surgical and restorative manipulation, demonstrating a lower tendency for recession but a higher risk for inflammatory enlargement and pocket formation if plaque control is inadequate) (Malpartida-Carrillo *et al.*, 2021).

The biotype fundamentally influences the tissue's response to the prosthetic interface. Patients with a thin biotype are at an elevated risk of "gingival retreat" following the placement of subgingival margins. Because the blood supply and connective tissue volume are limited, the inflammatory response to even minimal subgingival encroachment, or the trauma of impression taking, often results in permanent apical migration of the gingival margin (Abdulhady *et al.*, 2025).

In cases where the underlying tooth structure is discolored, the thin biotype provides minimal soft tissue opacity. Consequently, the ceramic material may appear gray or unnaturally bright through the thin gingival mucosa, necessitating a more aggressive subgingival preparation to mask the color transition, a decision that directly clashes with the biological need to preserve tissue health (Gadge *et al.*, 2025).

The thick phenotype offers greater stability, allowing for more predictable marginal positioning. However, the clinician

must be aware that in the event of inflammation, this biotype is more prone to pocketing rather than recession, which requires different periodontal management strategies (Damavandi *et al.*, 2026).

In thin biotypes, clinicians should prioritize supragingival or equigingival preparation margins to avoid the critical risk of post-operative recession. If subgingival placement is unavoidable, the restorative material must be carefully contoured to minimize pressure on the underlying soft tissue.

In cases where a thin biotype threatens the long-term stability of the aesthetic result, soft tissue grafting (such as connective tissue grafts) may be indicated prior to veneer preparation to "thicken" the tissue and provide a more stable biological base (Sarma *et al.*, 2022).

The patient must be informed about the limitations imposed by their periodontal phenotype. A realistic discussion regarding the stability of the gingival margin is essential, as the biological response is ultimately mediated by the patient's inherent tissue architecture.

The periodontal biotype is not merely an anatomical detail but a predictive tool for the biological response of the periodontium. By integrating this assessment into the diagnostic phase, the clinician can transition from a reactive approach to a proactive, personalized strategy that preserves tissue integrity while achieving superior aesthetic integration.

2.5. Management of Emergence Profile


The emergence profile represents the geometric transition between the natural root surface and the emergence of the prosthetic crown (the ceramic veneer) from the gingival sulcus. In the context of ceramic veneers, precise management of this contour is pivotal; it serves as the dynamic interface where the restorative material interacts with the marginal gingiva, directly influencing both esthetic harmony and the long-term biological response of the periodontium.

The profile of the veneer at the cervical level must be carefully engineered to respect the natural anatomy of the tooth (Granell-Ruiz *et al.*, 2023). A common clinical error is

the creation of over-contoured (convex) restorations. Such profiles force the gingival tissue to displace, creating a chronic inflammatory environment. An ideal emergence profile should follow a nearly linear or slightly concave transition from the finish line to the gingival margin, allowing for physiologic "creeping" or stable positioning of the marginal soft tissue.

If the restorative contour is too bulky, the resulting compression of the gingiva disrupts the blood supply, leading to recession. Conversely, an under-contoured profile, while less harmful to the gingiva, may create a "shelf" that encourages biofilm accumulation and food impaction.

The emergence profile is established at the intersection of three zones: the margin, the transmucosal zone, and the coronal contour. Any discrepancy (step or ledge) at the junction creates a stagnation site for plaque, which is impossible to clean effectively, inevitably leading to localized gingivitis. The transmucosal transition must mimic the natural convexity of the tooth. By maintaining a flat profile, the clinician facilitates proper mechanical plaque control by the patient's toothbrush and interdental cleaning aids (dental floss or interdental brushes), which is critical for the maintenance of the gingival seal (Fig. 1).



Achieving an idealized emergence profile requires meticulous collaboration between the clinician and the dental technician:

- Accurate impression and soft tissue management: the emergence profile cannot be created if the gingival sulcus is not adequately captured. The use of retraction cords or laser troughing is essential to expose the finish line and the immediate subgingival root surface, allowing the laboratory to accurately reproduce the root anatomy (Inchara *et al.*, 2022).

- "Digital" and "analog" communication: the use of diagnostic wax-ups or digital mock-ups serves as the blueprint for the emergence profile. The clinician must ensure that the transition from the prepared tooth to the veneer is not

dictated solely by material thickness but by the biological requirements of the existing gingival architecture (Moldovani *et al.*, 2021).

- The role of provisional restorations: for complex cases, temporary veneers act as a "training" tool for the gingival tissue. By adjusting the emergence profile of the temporaries and monitoring the gingival response over several weeks, the clinician can "sculpt" the soft tissue, ensuring the final ceramic veneer will be integrated harmoniously and naturally (Hammond *et al.*, 2022).

- Surface finishing and contouring: even if the prosthetic design is correct, improper polishing can create a "false" contour (Sasany *et al.*, 2022). Post-cementation adjustment must be performed

with extreme caution using fine-grit burs and polishing systems to ensure that the emergence profile remains smooth and consistent.

The ultimate goal of managing the emergence profile is the achievement of periodontal stability. A well-contoured veneer allows for the formation of a healthy sulcus, where the gingiva can remain at the optimal level relative to the cervical margin of the restoration. When the emergence profile correctly mimics the natural tooth anatomy, the gingival tissue exhibits better health, improved color, and superior architectural stability, all of which are essential to prevent the exposure of the veneer margin and the subsequent "aging" of the restoration.

2.6. The relationship between gingival zenith and symmetry

In the pursuit of an optimal esthetic outcome for ceramic veneer restorations, the gingival zenith (the most apical point of the gingival margin of a tooth) acts as the primary reference point for gingival architecture. Achieving bilateral and segmental symmetry is a critical requirement for creating a natural, harmonious smile. The manipulation of these zeniths, often required in veneer therapy, must be performed with a profound respect for periodontal anatomy to ensure that symmetry does not come at the cost of biological health.

The gingival zenith is not uniform across the dentition. Its position is dictated by the tooth anatomy and its location within the dental arch. For central incisors and canines, the zenith is typically located slightly distal to the long axis of the tooth. In contrast, for lateral incisors, the zenith is often centered. Symmetry in the smile arc is largely achieved when the zeniths of the contralateral teeth (right central incisor and left central incisor) are at the same vertical level and follow a harmonious transition, where the zeniths of the lateral incisors are positioned slightly more coronally than those of the central incisors and canines.

When designing veneers for the "esthetic zone," the clinician often faces discrepancies in the gingival levels due to previous trauma, periodontal disease, or altered passive

eruption. When symmetry is lacking, the clinician may need to perform a gingivectomy or a crown-lengthening procedure. This surgical intervention must carefully respect the biological width (Jurado *et al.*, 2024). Moving a zenith apically is restricted by the position of the alveolar bone; if the bone is not repositioned simultaneously, the gingival tissue will eventually migrate coronally, leading to a "relapse" of the gingival margin and loss of the newly created symmetry.

A common error is attempting to achieve symmetry by placing the veneer margin in a way that physically forces the gingival tissue to a new position. If the restoration imposes a permanent displacement on the soft tissue, it creates a constant source of pressure, leading to inflammation and, ultimately, uncontrolled recession that disrupts the very symmetry the clinician sought to create.

To ensure that the gingival zeniths contribute to a stable and beautiful result, the following clinical strategies are employed:

- Diagnostic mock-up as a guide: the desired gingival zenith position should be established during the diagnostic wax-up phase. This allows the clinician to determine if the required symmetry can be achieved through soft tissue surgery or if it requires restorative compensation (Guarnieri *et al.*, 2025).
- Conservative gingival correction: whenever possible, corrections should be limited to the gingival sulcus (gingivoplasty) if the underlying biological width allows. If the correction exceeds 1.0 - 1.5 mm, a periodontal evaluation for potential osseous resection is mandatory to avoid unpredictable tissue remodeling.
- Alignment with the smile arc: the zeniths must be viewed in the context of the overall smile arc (the relationship between the incisal edges and the lower lip). Achieving zenith symmetry while ignoring the alignment of the incisal edges will result in a visual disharmony, even if the gingival margins themselves are perfectly levelled (Freitas *et al.*, 2024).
- Integration with the emergence profile: the zenith is the "peak" of the emergence profile. Consequently, the transition from

the margin to the free gingival margin must be seamless at the zenith point. Any over-contouring at this specific location is highly visible and prone to trapping debris, as it is the most difficult area for a patient to reach during interdental cleaning.

The stability of the gingival zenith is the ultimate test of a successful veneer treatment. A zenith that migrates post-operatively, either due to biological width violation, poor emergence profile or surgical trauma, renders the entire esthetic effort a failure. By respecting the anatomical limits of the gingiva and utilizing precise, non-traumatic surgical and restorative techniques, the clinician can ensure that the symmetry achieved at the time of delivery remains stable over the long term.

2.7. Long-term monitoring

The delivery of ceramic veneers marks the beginning of the "maintenance phase," which is as critical to the longevity of the restoration as the clinical procedures themselves. Long-term monitoring is not merely a passive follow-up; it is an active diagnostic process designed to detect early signs of biological or mechanical failure. In the context of the periodontal-restorative interface, monitoring focuses on the dynamic stability of the soft tissues and the integrity of the marginal seal.

Post-operative evaluation must be systematic and quantitative to identify subtle changes before they become irreversible. Key parameters include: gingival margin stability (precise documentation of the gingival margin position relative to the ceramic finish line is mandatory; any sign of apical migration, recession, should trigger an investigation into the emergence profile, patient hygiene compliance, or potential biological width encroachment), probing depth and bleeding on probing - BOP (a gentle assessment of the sulcus is essential; the presence of persistent BOP, especially when localized to a single veneer margin, is a hallmark of subgingival biofilm accumulation or residual cement, both of which necessitate immediate professional intervention) and tissue architecture and texture (healthy peri-veneer tissue should appear pale pink, firm,

and stippled; any shift toward a "rolled," erythematous, or edematous appearance indicates a disruption in the biological seal, often related to an improper emergence profile that prevents effective plaque removal) (Sen *et al.*, 2024).

The success of long-term monitoring is heavily dependent on the patient's ability to maintain the peri-veneer environment. Patients must be educated on the specific challenges of cleaning veneer margins. This includes the use of soft-bristled brushes, non-abrasive toothpastes, and the appropriate use of interdental cleaning aids (Almutairi *et al.*, 2025). Patients with ceramic restorations should be enrolled in a periodic recall program (typically every 3 to 6 months). These visits allow for the professional cleaning of the veneer margins with specialized instruments (sonic or ultrasonic scalers with specialized tips) and the polishing of the ceramic surfaces to maintain a low-roughness profile.

Long-term monitoring allows the clinician to identify complications that may only become apparent years after placement. Over time, the luting agent may undergo hydrolytic degradation. Monitoring for discoloration at the margin or "staining" of the interface can indicate the loss of the peripheral seal, which may predispose the tooth to secondary caries. As patients age, the gingival zenith may naturally shift or the soft tissue may undergo minor recession. The clinician must differentiate between physiological tissue changes and pathological responses to the veneer, adjusting the home care routine accordingly (Mihalie *et al.*, 2022). Periodic checks of the occlusal scheme are vital. Excess force on the veneer can lead to stress concentration at the cervical margin, which may cause micro-fractures in the ceramic or, in extreme cases, trigger inflammatory periodontal reactions due to tooth mobility (Ladino *et al.*, 2022).

Standardized clinical photography and digital records are the most effective tools for long-term monitoring. By comparing current images with baseline clinical records taken at the time of cementation, the clinician can objectively assess the stability of the gingival

zenith, the emergence profile, and the overall periodontal phenotype (Soares-Rusu *et al.*, 2024). This objective data allows for timely,

evidence-based adjustments, effectively extending the lifespan of the restoration.

4. CONCLUSIONS

Based on the analysis of current periodontal and restorative protocols, the following conclusions can be drawn:

1. The long-term stability of ceramic veneers is fundamentally dependent on the respect for the biological width. Clinical procedures must prioritize the preservation of the junctional epithelium and supracrestal connective tissue attachment to prevent chronic inflammation and subsequent recession.
2. The emergence profile serves as the primary dynamic interface between the restoration and the periodontium. A flat, non-bulky contour is essential to facilitate patient-led plaque control and to prevent iatrogenic pressure on the soft tissues, which is a leading cause of gingival pathology.
3. The surface quality of the ceramic material is a determinant factor in cellular adhesion. Polishing protocols must ensure a surface roughness (Ra) below 0.2 μm to promote a stable hemidesmosomal attachment and to minimize biofilm accumulation.
4. Residual cement acts as a major catalyst for periodontal breakdown. The implementation of rigorous, magnified inspection protocols and precise cementation techniques is non-negotiable to avoid the inflammatory cascade associated with subgingival cement remnants.
5. The periodontal biotype acts as a prognostic marker for the biological response. Thin-scalloped biotypes require a more conservative, often supragingival approach to margin placement, whereas thick-flat biotypes offer more flexibility but carry specific risks of pocketing under suboptimal hygiene conditions.
6. The aesthetic harmony achieved through the positioning of the gingival zenith must be balanced with the anatomical limits of the alveolar bone. Surgical or restorative interventions aimed at zenith realignment must respect the biological limits of the periodontium to ensure permanent and predictable stability.

REFERENCES

- 1 Abdulhady E.M., Soliman O.H., Al Bahrawy M. Stability of Gingival Margin Subsequent Esthetic Crown Lengthening in Thick Periodontal Biotype versus Thin Periodontal Biotype; Prospective Clinical Trial. *Egyptian Dental Journal*; 2025;71(2), 1279-1292.
- 2 Alharbi E.A., Najar A.A., Daghriri F.A., Ali A.M. Biocompatible Issues Related to Restorative Dental Material-II (Adverse Reactions). *Journal of Advanced Medical and Dental Sciences Research*; 2024;12(3), 12-20.
- 3 Almutairi D., Alshahrani S., Balawi A., Alnasser S., Alshamlan A., Almugim H., Alharbi M.A. Level of Awareness About Dental Veneers and Their Oral Hygiene Care Among the General Saudi Public: A Cross-Sectional Study. *Healthcare*; 2025;13, 2170.
- 4 Al-Shorman H.M., Abu-Naba'a L.A., Sghaireen M.G., Alam M.K. The Effect of Various Preparation and Cementation Techniques of Dental Veneers on Periodontal Status: A Systematic Review and Meta-Analysis. *European Journal of Dentistry*; 2024;18(2):458-467.
- 5 Andrei P.V., Mihaela S.S., Andreea T., Cosmin C., Andrian N., Norina F. THE PROVISIONAL TRANSITIONAL PROSTHETIC REHABILITATION OF EDENTULOUS PATIENTS WITH PERIODONTAL PATHOLOGY. *Romanian Journal of Medical and Dental Education*; 2024; 13(6).
- 6 Ceravolo M., Castro F., González-Mosquera A., López-Solache A., Manarte-Monteiro P., Pereira da Silva L. Adhesive Restoration Performance in Deep Subgingival Margins: Deep Margin Elevation

- Versus Surgical Crown Lengthening—A Systematic Review. *Adhesives*; 2026: 2(2), 9.
- 7 Cortés-Sánchez M.V., Villa-Bedoya E.I., Villegas-Vélez M., Botero JE. The Influence of Full-Crown Preparation Margin Position on Periodontal Outcomes: A Systematic Review and Meta-Analysis. *Current Oral Health Reports*; 2026:13(1), 10.
- 8 Damavandi M., Azizian R., Shojaei N. Evaluation of gingival biotype and bone thickness in maxillary implant patients: A clinical study based on cone-beam computed tomography. *Journal of Advanced Periodontology & Implant Dentistry*; 2026: 18(2), 90-97.
- 9 De Angelis N., Pesce P., Caponio V.C.A., Santamaria G., Spanu O., Menini M. Marginal Bone Level Changes in Full-Arch Rehabilitation: Digital Versus Analog Protocols-A 5-Year Retrospective Study. *Clinical Implant Dentistry and Related Research*; 2025:27(4):e70080.
- 10 Ercoli C., Tarnow D., Poggio C.E., Tsigarida A., Ferrari M., Caton J.G., Chochlidakis K. (2021). The relationships between tooth-supported fixed dental prostheses and restorations and the periodontium. *Journal of Prosthodontics*; 2021:30(4), 305-317.
- 11 Freitas K. Evaluation of the attractiveness of different gingival zeniths in smile esthetics. *Dental press journal of orthodontics*; 2024.
- 12 Gadge N.P., Chawla R., Ronad S., Bhole S.V., Kumar P., Kattimani P.T. Correlations of gingival biotype with clinical crown and periodontal parameters in maxillary and mandibular jaws. *Clinical Advances in Periodontics*; 2025:15(1), 77-85.
- 13 Gargiulo W., Wentz M., Orban b. Dimensions and relations of the dentogingival junction in humans. *Journal of Periodontology*; 1961:32: 261-267. 19.
- 14 Granell-Ruiz M., Rech-Ortega C., Oteiza-Galdón B., Bouazza-Juanes K. Case report: Vertical preparation protocol for veneers. *Journal of Clinical and Experimental Dentistry*; 2023: 15(4), e346.
- 15 Guarnieri F.D.F., Perazza B., Briso A.L.F., Ramos F.D.S., Esteves L.M.B., Oliveira L.D.C., Fagundes T.C. Short-term aesthetic treatment with gingival recontouring and anterior veneers using a full digital workflow. *Revista Odontologica Araçatuba*; 2025:37-42.
- 16 Hammond B.D., Machowski M., Londono J., Pannu D. Fabrication of Porcelain Veneer Provisional Restorations: A Critical Review. *Dentistry Review*; 2022:2(2), 100004.
- 17 Inchara R., Karthickraj S.M. Evaluation of finish line exposure with retraction cord placement in esthetic procedures-a retrospective study. *Journal of Research in Medical and Dental Science*; 2022:10.
- 18 Jurado C.A., Villalobos-Tinoco J., Lackey M.A., Rojas-Rueda S., Robles M., Tsujimoto A. Three dimensional-printed gingivectomy and tooth reduction guides prior ceramic restorations: A case report. *Dentistry journal*; 2024:12(8), 245.
- 19 Karvekar S., Aathilingaperumal S., Sarada V. Periodontal Tissue Response to Crown Margin Placement: A Comparative Evaluation of Prosthodontic Designs. *European Journal of Prosthodontics & Restorative Dentistry*; 2025:33, 465.
- 20 Ladino L., Sanjuan M.E., Valdéz D.J., Eslava R.A. Clinical and biomechanical performance of occlusal veneers: a scoping review. *The Journal of Contemporary Dental Practice*; 2022:22, 1327-1337.
- 21 Malpartida-Carrillo V., Tinedo-Lopez P.L., Guerrero M.E., Amaya-Pajares S.P., Özcan M., Rösing, C.K. Periodontal phenotype: A review of historical and current classifications evaluating different methods and characteristics. *Journal of Esthetic and Restorative Dentistry*; 2021:33(3), 432-445.
- 22 Metwally NM, Elshenawy E.A., Elghareb L.A. Surface roughness and wear performance of Bioflx versus stainless-steel primary crowns (an in-vitro study). *BMC Oral Health*; 2025:25(1), 343.
- 23 Mihali S.G., Lolos D., Popa G., Tudor A., Bratu D.C. Retrospective long-term clinical outcome of feldspathic ceramic veneers. *Materials*; 2022:15(6), 2150.
- 24 Moldovani D., Diamantopoulou S., Edelhoff D., Papazoglou E. Accuracy of Indirect Veneer Mock-Up in Comparison to Diagnostic Wax-Up. *International Journal of Prosthodontics*; 2023: 36(4).
- 25 Montevecchi M., Valeriani L., Salvadori M.F., Stefanini M., Zucchelli G. Excess cement and peri-implant disease: A cross-sectional clinical endoscopic study. *Journal of Periodontology*; 2025: 96(9), 965-973.
- 26 Nenov S., Iliev G. Emergence profile of porcelain veneers and gingival design in the digital workflow. *ПРОБЛЕМИ НА ДЕНТАЛНАТА МЕДИЦИНА*; 2025: 55.
- 27 Niemczyk W., Niemczyk S., Prokurat M., Grudnik K.E., Migas M., Wągrowiska K., Kasperczyk J

- Etiology of gingival recession-a literature review. *Wiadomości Lekarskie*; 2024;77(5), 1080-1085.
- 28 Parihar A.S., Nandanan N., Kurien V.T.V., Bandgar S., Patri G., Laddha R. Impact of Periodontal Biotype on the Long-Term Esthetics of Implant-Supported Restorations: A 5-Year Prospective Study. *Journal of Pharmacy and Bioallied Sciences*; 2025;17(Suppl 1), S555-S557.
 - 29 Paul S., Hanisch O., Nesic D. Human gingival fibroblast proliferation on materials used for dental implant abutments: a systematic review. *International Journal of Prosthodontics*; 2021;34(6).
 - 30 Payaminia L., Moslemian N., Younespour S., Koulivand S., Alikhasi M. Evaluating the effect of repeated use of milling burs on surface roughness and adaptation of digitally fabricated ceramic veneers. *Heliyon*; 2021;7(4).
 - 31 Sarma M., Shenoy N., Bhandary R. Gingival biotype: a secret for esthetic success. *Journal of Health and Allied Sciences NU*; 2022;12(01), 13-17.
 - 32 Sasany R., Ergün Kunt G., Koca M.F. Influence different polishing systems on roughness and colour stability of chairside CAD/CAM blocks with laminate veneer thickness. *Journal of Applied Biomaterials & Functional Materials*; 2022;20, 22808000221086493.
 - 33 Sen N., Olley R.C. Retrospective evaluation of factors affecting long-term clinical performance of CAD-CAM laminate veneers. *International Journal of Prosthodontics*; 2024; 37(4), 411-416.
 - 34 Skevi L., Reeksting B.J., Hoffmann T.D., Gebhard S., Paine K. Incorporation of bacteria in concrete: the case against MICP as a means for strength improvement. *Cement and Concrete Composites*; 2021;120, 104056.
 - 35 Soares-Rusu I.B., Villavicencio-Espinoza C.A., de Oliveira N.A., Wang L., Honório H.M., Rubo J. H., Borges A.F. Using digital photographs as a tool to assess the clinical color stability of lithium disilicate veneers: A clinical trial. *The Journal of Prosthetic Dentistry*; 2024; 131(5), 859-864.
 - 36 Srimaneepong V., Heboyan A., Zafar M.S., Khurshid Z., Marya A., Fernandes G.V., Rokaya D. Fixed prosthetic restorations and periodontal health: a narrative review. *Journal of functional biomaterials*; 2022; 13(1), 15.
 - 37 Talpos S., Titihazan F., Veja I., Dobos M., Rominu M., Negrutiu M.L., Hajaj T. The Role of Preparation Geometry in Ceramic Veneers: A Comprehensive Literature Review. *Medicine in Evolution*; 2025; 31(2), 120-134.
 - 38 Yinuo Y., Qiqian Y., Kexin C., Zhihan W., Qiulan L., Ke Z., Yutao J., Xiaoshi J. Silk fibroin microgrooved zirconia surfaces improve connective tissue sealing through mediating glycolysis of fibroblasts. *Materials Today Bio*; 2024;27,101158.
 - 39 Zihan S., Xiaoqian Y., Qingxian L. Emergence Profile-specific Biological Width and Zonal Mucosal Thickness: A Digital Analysis of the Peri-implant Soft Tissue Phenotype and Its Clinical Relevance. *Research Square*; 2025.